

2015

## The Influence of Short-term Quercetin Supplementation on Peak Oxygen Uptake during Simulated Altitude Exposure in Trained Cyclists

Justin Carlstrom

Jamie M. Vener

Justin L. Matthews

*California State University, Monterey Bay, [jmatthews@csumb.edu](mailto:jmatthews@csumb.edu)*

Ashleigh Raub

Follow this and additional works at: [https://digitalcommons.csumb.edu/psy\\_fac](https://digitalcommons.csumb.edu/psy_fac)

---

### Recommended Citation

Carlstrom, J., Vener, J.M., & Matthews, J.L., & Raub, A. (2015). The influence of short-term quercetin supplementation on peak oxygen uptake during simulated altitude exposure in trained cyclists. *International Journal of Exercise Science*, 8, 394-402.

This Article is brought to you for free and open access by the Psychology at Digital Commons @ CSUMB. It has been accepted for inclusion in Psychology Faculty Publications and Presentations by an authorized administrator of Digital Commons @ CSUMB. For more information, please contact [digitalcommons@csumb.edu](mailto:digitalcommons@csumb.edu).

---

## The Influence of Short-term Quercetin Supplementation on Peak Oxygen Uptake during Simulated Altitude Exposure in Trained Cyclists

JUSTIN CARLSTROM<sup>#1</sup>, JAMIE M. VENER<sup>#2</sup>, JUSTIN L. MATTHEWS<sup>#3</sup>, and ASHLEIGH RAUB<sup>\*1</sup>

<sup>1</sup>Castleton College, Castleton, VT, USA; <sup>2</sup>Southern Oregon University, Ashland, OR, USA; <sup>3</sup>California State University, Monterey Bay, Seaside, CA, USA

‡Denotes professional author, \*Denotes undergraduate student author

---

### ABSTRACT

*International Journal of Exercise Science 8(4): 394-402, 2015.* Endurance performance and peak oxygen uptake ( $VO_{2peak}$ ) decline as altitude increases. Some data exist suggesting that quercetin supplementation improves aerobic capacity in trained and untrained individuals at sea-level (normobaric normoxic conditions). Few studies have examined the effects of quercetin on endurance performance during simulated altitude exposure (normobaric hypoxic conditions). The purpose of this study was to determine the effects of consuming  $1000 \text{ mg} \cdot \text{day}^{-1}$  of quercetin for two weeks on cycling  $VO_{2peak}$  in healthy trained male cyclists performing under normobaric normoxic and hypoxic conditions (NP and HP, respectively). Fourteen age-matched healthy male subjects were randomized to either a placebo or quercetin group. Baseline and post supplementation  $VO_{2peak}$  values were quantified during incremental cycling under normobaric normoxic ( $FIO_2 = 20.9\%$ ) and normobaric hypoxic ( $FIO_2 = 13.6 \pm 0.2\%$ ) conditions. Subjects consumed capsules twice daily with either 500 mg quercetin or placebo (Tang) for two weeks and were re-assessed. Test order was randomized and assessments were separated by 48-72 hours. At baseline, there were no significant differences between groups for  $VO_{2peak}$  normobaric normoxic trials (NP<sub>baseline</sub> Placebo vs. NP<sub>baseline</sub> Quercetin =  $58.7 \pm 8.8$  and  $61.5 \pm 7.9 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $p = 0.541$ ) and normobaric hypoxic trials (HP<sub>baseline</sub> Placebo vs. HP<sub>baseline</sub> Quercetin =  $48.5 \pm 8.3$  and  $50.8 \pm 4.8 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $p = 0.534$ ). No significant differences were found after treatment (Placebo: NP<sub>baseline</sub> vs. NP<sub>post</sub> =  $58.7 \pm 8.8$  and  $56.7 \pm 7.4 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $p = 0.10$ ; HP<sub>baseline</sub> vs. HP<sub>post</sub> =  $48.5 \pm 8.3$  and  $47.1 \pm 8.3$ , respectively,  $p = 0.50$ ; Quercetin: NP<sub>baseline</sub> vs. NP<sub>post</sub> =  $61.5 \pm 7.9$  and  $62.4 \pm 7.3 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $p = 0.558$ ; HP<sub>baseline</sub> vs. HP<sub>post</sub> =  $50.8 \pm 4.8$  and  $51.2 \pm 3.8 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $p = 0.656$ ). These data suggest that short-term quercetin supplementation at  $1000 \text{ mg} \cdot \text{day}^{-1}$  does not affect  $VO_{2peak}$  elicited via incremental maximal cycle testing under normobaric normoxic and hypoxic conditions in trained male cyclists. Furthermore, quercetin supplementation did not attenuate the decline in  $VO_{2peak}$  that was evident in the normobaric hypoxic condition.

KEY WORDS: Hypoxia,  $VO_{2Peak}$ , cycling, endurance performance

### INTRODUCTION

In recent years, much attention has been given to the role of dietary antioxidants in

preventing disease and improving overall health. Quercetin, a polyphenolic flavonoid, is a naturally occurring substance that is found in a variety of food.

Common food sources of quercetin include tea, onions, apples, peppers, blueberries and dark green vegetables, with the average intake of U.S. adults at approximately 4 mg · day<sup>-1</sup>. However, daily intakes may be as high as 200 – 500 mg in individuals consuming a diet high in fruits and vegetables (5). Although not entirely conclusive, epidemiologic studies provide evidence that self-selected diets high in quercetin may play a role in preventing various diseases. High quercetin intake has been linked to a reduced risk of colorectal, kidney, pancreatic, prostate, and lung cancer, as well as cardiovascular disease (10, 11, 29). *In vitro*, animal, and human studies have shown that quercetin possesses antioxidant, anti-carcinogenic, anti-inflammatory, and cardioprotective properties (3, 12, 14, 23, 26). In addition, there is evidence that quercetin may lower blood pressure through vasodilatory effects (14, 19). Based on these studies, quercetin appears to have significant health benefits.

Because of the purported health benefits of quercetin, investigators have begun to examine the efficacy of quercetin in improving athletic performance. Regular aerobic exercise can increase muscle mitochondrial density by 20% to 100%, depending on the workload (13). This increase in mitochondrial density contributes to improved fitness and performance by increasing the aerobic capacity of skeletal muscle. Limited data show that quercetin may have the ability to induce mitochondrial biogenesis and improve aerobic exercise performance. Human studies have shown that daily supplementation of quercetin (500 mg twice daily) improves aerobic capacity, time to exhaustion, time trial performance, and

exercise tolerance (20, 21). Furthermore, recent evidence from both animal and human studies suggests that quercetin supplementation resulted in increased mitochondrial capacity, oxidative enzyme activity, and increased time to exhaustion (8, 9). Additional data show that quercetin may cause vasodilation, resulting in improved blood flow (5, 18, 22). An increase in blood flow to working muscle, coupled with an increase in mitochondrial density, could result in improved endurance performance by increasing an individual's aerobic capacity and fat utilization. Although these data support the use of quercetin to improve aerobic performance, other studies have shown that quercetin has no effect on aerobic capacity (1, 6). More work is needed in this area to better define the efficacy of quercetin in improving athletic performance under various conditions.

It is well established that altitude decreases VO<sub>2peak</sub> in humans and evidence suggests that oxidative stress increases at altitude (16, 27). Consequently, the use of nutritional interventions, such as antioxidant supplementation to enhance antioxidant capacity and prevent and/or treat symptoms of acute mountain sickness has gained attention. Although justification for their use is not based on scientific evidence, *Ginkgo Biloba* and *Hippophae rhamnoides* L. have been used for centuries in Chinese and Tibetan medicine to combat AMS (24). Data from Purushothama et al. (2008) show that *H. rhamnoides* exerted a protective effect in rats exposed to hypoxia (24). Quercetin is a major component of this supplement (30). However, there is very limited data regarding the efficacy of isolated quercetin in attenuating the effects

of hypoxia *in vivo*. Zhou et al. investigated the protective effect of quercetin in rats exposed to a chronic (23 h per day) hypobaric, hypoxic environment (31). Quercetin attenuated the decline in blood pH, PO<sub>2</sub>, SpO<sub>2</sub>, and PCO<sub>2</sub> that occurred in the control group. In addition, quercetin increased nitric oxide levels, a potent vasodilator (31). Based on this limited data, it is plausible that quercetin may exert protective effects in an individual exposed to a hypoxic environment.

Significant reductions in VO<sub>2peak</sub> generally occur at an altitude of 1,500 to 1,600 m (5,000 ft). Beyond this altitude, VO<sub>2peak</sub> decreases approximately 8% to 11% for every 1,000 m increase (or 3% for every 1,000 ft increase) (27). In addition, variables such as heart rate, ventilation, and perception of fatigue, at rest and during exercise, are increased at higher elevation. The lack of oxygen at altitude also affects substrate metabolism. At altitude, there is an increased reliance carbohydrate for fuel since this substrate can be metabolized anaerobically (4, 17). Therefore, aerobic exercise at altitude may be limited, in part, by glycogen depletion, as this is a very limited fuel supply. Due to the detrimental effect of altitude on athletic performance, many studies have investigated various nutritional strategies in an effort to offset the performance decrement typically seen in an individual performing in that environment. Recent studies have examined the efficacy of antioxidants and other nutritional supplements in 1) attenuating the decline in VO<sub>2max</sub> and 2) enhancing fat utilization, thus sparing glycogen, in individuals exercising at altitude. However, there is very limited data regarding the efficacy of quercetin in

enhancing aerobic performance and fat utilization at altitude. The overall goal of this investigation is to examine the effect of quercetin on peak oxygen uptake (VO<sub>2peak</sub>) in trained individuals exercising under normobaric normoxic and hypoxic conditions.

## METHODS

### *Participants*

Fourteen healthy male subjects (age: 38.9 ± 9.8 y, weight: 75.7 ± 7.9 kg) were recruited through flyers posted on the Castleton State College campus and throughout the Rutland community, and through solicitation of the Killington Pico Cycling Club. Prior to data collection, participants were advised of the possible risks, methods, and potential benefits of participation in the study. A health history questionnaire and informed consent (approved by the Castleton State College Institutional Review Board for research involving human subjects) was completed by each subject.

### *Protocol*

Subjects were age-matched and randomized to either a placebo group (consumed one capsule twice daily with 500 mg Tang from the Kraft Foods Company, Northfield, IL) or quercetin supplementation group (consumed one capsule with 500 mg twice daily with quercetin from Jarrow Formulas, Los Angeles, CA) in a double-blind design. Subjects had no significant recent exposure to altitude training or racing and were asked to maintain their typical training and dietary routines for the duration of the study. Subjects were instructed to refrain from strenuous exercise and excessive alcohol for 24 hours prior to testing.

Subjects maintained a dietary record for 48 hours prior to initial testing and were asked to match it as closely as possible for each successive trial. All maximal trials were separated by 48-72 hours.

Metabolic data was collected using a ParvoMedics metabolic system (ParvoMedics, Salt Lake City, UT). All testing was conducted using an electronically-braked cycle ergometer designed to interface with the subject's personal cycle of choice (Velotron DynaFit Pro, Seattle, WA). Subjects completed a five minute warm-up at a workload of 50 watts at the start of each trial. Using a continuous incremental protocol, workload increased by 25 watts per minute until volitional exhaustion was achieved. All tests were terminated after subjects failed to maintain their final workload for 15 seconds and/or voluntarily identified that their maximal effort was achieved.  $VO_{2peak}$  was identified as the highest value recorded during the final workload achieved by each subject.

Subjects performed cycling trials under normobaric normoxic conditions ( $FIO_2 = 20.9\%$ ,  $752 \pm 4$  mmHg) and hypoxic conditions ( $FIO_2 = 13.6 \pm 0.2\%$ ,  $754 \pm 3$  mmHg; simulated approximately 3600 meters above sea-level). Baseline and post treatment normoxic and hypoxic trials were randomized and subjects were not informed of the trial condition. Hypoxic conditions were produced using a hypoxic generator (Colorado Altitude Training, Louisville, CO) connected to two, 200 L Douglas bags that were used to create a reservoir of inspiratory gas for the subjects during hypoxic trials (Figure 1). In order to avoid subject bias, this method was also used during the normoxic trials with the Douglas bags filled with 20.9%  $O_2$ .



Figure 1. Experimental set up.

### Statistical Analysis

A MANOVA examined oxygen level (normoxia/hypoxia), quercetin supplementation, and time as independent variables and  $VO_{2peak}$  and power (w) as dependent variables (Wilks'  $\lambda = .095$ ,  $F(9,4) = 4.24$ ,  $p = .09$ ). Univariate ANOVAs examined differences in individual dependent variables. Thereafter, post-hoc t-tests were used to examine pair-wise differences. Using Bonferroni correction, a family-wise alpha of .05 was maintained for all post-hoc pair-wise comparisons.

## RESULTS

$VO_{2peak}$  was not different between groups at baseline for normobaric normoxic trials (placebo vs. quercetin group =  $58.7 \pm 8.8$  and  $61.5 \pm 7.9$   $ml \cdot kg^{-1} \cdot min^{-1}$ , respectively,  $t(12) = -.63$ ,  $p = 0.54$ ) (Table 1) and hypoxic trials (placebo vs. quercetin group =  $48.5 \pm 8.3$  and  $50.8 \pm 4.8$   $ml \cdot kg^{-1} \cdot min^{-1}$ , respectively,  $t(12) = -.64$ ,  $p = 0.53$ ) (Table 2). There was no statistically significant difference found in  $VO_{2peak}$  during normoxic trials for the placebo group when comparing baseline values to post placebo supplementation values ( $58.7 \pm 8.8$  and  $56.7 \pm 7.4$   $ml \cdot kg^{-1} \cdot min^{-1}$ , respectively,  $t(6) = 1.95$ ,  $p = 0.10$ ) (Table 1). There was no statistically

significant difference found in  $VO_{2peak}$  during hypoxic trials for the placebo group when comparing baseline values to post placebo supplementation values ( $48.5 \pm 8.3$  and  $47.1 \pm 8.3 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $t(6) = .72, p = 0.50$  (Table 2). There was no statistically significant difference found in  $VO_{2peak}$  during normoxic trials for the quercetin supplementation group when comparing baseline values to post supplementation values ( $61.5 \pm 7.9$  and  $62.4 \pm 7.3 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $t(6) = -.62, p = 0.56$ ) (Table 1). There was no statistically significant difference found in  $VO_{2peak}$  during hypoxic trials for the quercetin supplementation group when comparing baseline values to post supplementation values ( $50.8 \pm 4.8$  and  $51.2 \pm 3.8 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , respectively,  $t(6) = -.47, p = 0.66$ ) (Table 2). As expected,  $VO_{2peak}$  for all hypoxic trials was significantly ( $p < 0.05$ ) reduced when compared with normoxic trials (baseline normoxia vs. baseline hypoxia and post normoxia vs. post hypoxia) trails for all groups (range = -16.9% to -17.9% of baseline  $VO_{2peak}$ ) (Table 3). These data suggest that short-term quercetin supplementation for 2 weeks at  $1000 \text{ mg} \cdot \text{day}^{-1}$  does not improve  $VO_{2peak}$  in healthy trained male cyclists performing under normobaric normoxic and hypoxic laboratory conditions.

**Table 1.** Mean ( $\pm$  SD) results for  $VO_{2peak}$  during normoxic trials for placebo and quercetin supplementation groups.

Normoxic Trials	Peak $VO_2$ ( $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ )
Placebo baseline	$58.7 \pm 8.8^*$
Placebo post	$56.7 \pm 7.4^*$
Quercetin baseline	$61.5 \pm 7.9^*$
Quercetin post	$62.4 \pm 7.3^*$

**Table 2.** Mean ( $\pm$  SD) results for  $VO_{2peak}$  during hypoxic trials for placebo and quercetin supplementation groups.

Hypoxic Trials	Peak $VO_2$ ( $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ )
Placebo baseline	$48.5 \pm 8.3^{**}$
Placebo post	$47.1 \pm 8.3^{**}$
Quercetin baseline	$50.8 \pm 4.8^{**}$
Quercetin post	$51.2 \pm 3.8^{**}$

**Table 3.** Mean ( $\pm$  SD) results for  $VO_{2peak}$  comparing normoxic and hypoxic trials for placebo and quercetin supplementation groups.

Normoxic vs. Hypoxic Trials	Peak $VO_2$ ( $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ )		
	Normoxic	Hypoxic	% Decline
Placebo baseline	$58.7 \pm 8.8$	$48.5 \pm 8.3$	17.4 <sup>†</sup>
Placebo post	$56.7 \pm 7.4$	$47.1 \pm 8.3$	16.9 <sup>†</sup>
Quercetin baseline	$61.5 \pm 7.9$	$50.8 \pm 4.8$	17.4 <sup>†</sup>
Quercetin post	$62.4 \pm 7.3$	$51.2 \pm 3.8$	17.9 <sup>†</sup>

<sup>†</sup> $p < 0.0125$  (pair-wise)

## DISCUSSION

The primary purpose of this investigation was to evaluate the effects of quercetin supplementation on  $VO_{2peak}$  during incremental cycle ergometry under normobaric hypoxic conditions that simulated approximately 3600 meters of elevation above sea-level. In addition, the effect of quercetin supplementation on  $VO_{2peak}$  during normobaric normoxic/sea-level conditions was investigated. We report no beneficial effects of  $1000 \text{ mg} \cdot \text{day}^{-1}$  of quercetin supplementation for two weeks on  $VO_{2peak}$  under both hypoxic and normoxic conditions. While  $VO_{2peak}$  responses to normobaric hypoxia after quercetin supplementation have not been studied, our findings are consistent with others who have reported no beneficial effects to endurance performance and aerobic capacity after short-term (i.e., 2 - 4

weeks) quercetin supplementation (3, 4). Our findings are contrary to those who reported that quercetin supplementation improved cycling efficiency, aerobic capacity, fat utilization, and time trial performance in both trained and untrained individuals (7, 8, 9, 21).

Due to controversial findings related to the effects of quercetin supplementation on factors related to endurance performance in humans and animals, researchers have addressed the methodological variability of several of these studies. Jin and colleagues (2010) have questioned the bioavailability of extracted/processed versus natural forms of quercetin, suggesting that natural forms may be superior to processed forms. We chose a processed form for this investigation as dosage was easily controlled. Dosage has also varied from study to study. The 1000 mg · day<sup>-1</sup> used here is among the higher doses studied, although even higher doses may be needed to yield beneficial effects (15). It is possible that longer duration supplementation (i.e., > 4 weeks) and/or higher doses may be necessary to induce structural and functional biological changes that could enhance endurance performance or VO<sub>2peak</sub> under hypoxic conditions.

To our knowledge, the influence of quercetin supplementation on VO<sub>2peak</sub> during simulated altitude using normobaric hypoxia has not been investigated. Subudhi et al. (25) studied the effects of a combination of antioxidants on ventilatory threshold at altitude and reported that 30 days of supplementation improved acute ventilatory kinetics associated with adaptive responses to hypobaric hypoxia (4300meters), but there appeared to be no

long term benefits associated with the supplement. These researchers reported a 28% reduction in VO<sub>2peak</sub> at 4300 meters in both the control and treatment groups. Similarly, we found that VO<sub>2peak</sub> decreased by approximately 17% in both the placebo and quercetin supplementation group under normobaric hypoxic condition studied here (approximately 3600m simulated altitude), however there were no significant differences in VO<sub>2peak</sub> associated with quercetin supplementation.

Given that some studies have documented increased mitochondrial RNA, increased muscle blood flow and decreased blood pressure in response to quercetin supplementation, we felt that quercetin supplementation could be particularly beneficial to endurance athletes exercising at altitude. Our findings indicate no beneficial effects of short-term quercetin supplementation on VO<sub>2peak</sub> during maximal cycle ergometry testing at a simulated altitude of approximately 3600 meters. The dose used in this investigation was determined based on evidence from previous research reporting enhanced endurance performance after 1-4 weeks of quercetin supplementation at similar doses to those used here (20, 21). Human studies have shown that daily supplementation of quercetin (500 mg twice daily) improves aerobic capacity (VO<sub>2max</sub>), time to exhaustion, time trial performance, and exercise tolerance (20, 21). Furthermore, recent evidence from both animal and human studies suggests that quercetin supplementation resulted in increased mitochondrial capacity, oxidative enzyme activity, and increased time to exhaustion (8, 9). Additional data show that quercetin may cause vasodilation, resulting in

improved blood flow (18, 21, 22). An increase in blood flow to working muscle, coupled with an increase in mitochondrial density, could result in improved endurance performance by increasing an individual's aerobic capacity and fat utilization. Although these data support the use of quercetin to improve aerobic performance, our findings support those of others who have shown that quercetin has no effect on aerobic capacity as indicated by  $VO_{2peak}$  (1,6).

In conclusion, potential beneficial effects of quercetin supplementation that may be particularly relevant to endurance performance at altitude may involve structural changes at the muscle cell and vascular level (i.e., increased mitochondrial density, improved vasodilatory responses), therefore, longer durations of supplementation should be studied. Further, bioavailability of various forms of quercetin should be considered and natural forms of quercetin supplementation should be evaluated. Another area for study is the effects of quercetin supplementation and dose-response kinetics on untrained individual performing at altitude as these individuals may benefit more significantly from potential upregulation of structural components supporting  $VO_{2peak}$  and endurance performance at altitude.

#### ACKNOWLEDGEMENTS

Research reported in this publication was supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P20GM103449. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of NIGMS or NIH.

#### REFERENCES

1. Abbey EL, Rankin JW. Effect of quercetin supplementation on repeated-sprint performance, xanthine oxidase activity, and inflammation. *Int J Sport Nutr Exerc Metab* 21(2): 91-96, 2011.
2. Basnyat B, Cumbo TA, Edelman R. Acute medical problems in the Himalayas outside the setting of altitude sickness. *High Altitude Med Biol* 1(3): 167-174, 2001.
3. Boomgaarden I, Egert S, Rimbach G, Wolfram S, Muller MJ, Doring F. Quercetin supplementation and its effect on human monocyte gene expression profiles in vivo. *Br J Nutr* 104(3): 336-345, 2010.
4. Braun B. Effects of high altitude on substrate use and metabolic economy: cause and effect? *Med Sci Sports Exerc* 40(8): 1495-1500, 2008.
5. Chun OK, Chung SJ, Song WO. Estimated dietary flavonoid intake and major food sources of U.S. adults. *J Nutr* 137(5): 1244-1252, 2007.
6. Cureton KJ, Tomporowski PD, Singhal A, Pasley JD, Bigelman KA, Lambourne K. Dietary quercetin supplementation is not ergogenic in untrained men. *J Appl Physiol* 107(4): 1095-1104, 2009.
7. Davis JM, Carlstedt CJ, Chen S, Carmichael MD, Murphy EA. The dietary flavonoid quercetin increases  $VO_{2max}$  and endurance capacity. *Int J Sport Nutr Exerc Metab* 20(1): 56-62, 2010.
8. Davis JM, Murphy EA, Carmichael MD, Davis B. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Reg Integrat Compar Physiol* 296(4): R1071-1077, 2009.
9. Dumke CL, Nieman DC, Utter AC, Rigby MD, Quindry JC, Triplett NT. Quercetin's effect on cycling efficiency and substrate utilization. *Appl Physiol Nutr Metab* 34(6): 993-1000, 2009.
10. Egert S, Bosy-Westphal A, Seiberl J, Kurbitz C, Settler U, Plachta-Danielzik S. Quercetin reduces systolic blood pressure and plasma oxidised low-



density lipoprotein concentrations in overweight subjects with a high-cardiovascular disease risk phenotype: a double-blinded, placebo-controlled cross-over study. *Br J Nutr* 102(7): 1065-1074, 2009.

11. Ferris-Tortajada J, Berbel-Tornero O, Garcia-Castell J, Ortega-Garcia JA, Lopez-Andreu JA. Dietetic factors associated with prostate cancer. Protective effects of mediterranean diet. *Actas Urologicas Espanolas* 36(4): 239-245, 2012

12. Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of in vivo toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem Toxicol* 45(11): 2179-2205, 2007.

13. Hoppeler H, Fluck M. Plasticity of skeletal muscle mitochondria: structure and function. *Med Sci Sports Exerc* 35(1): 95-104, 2003.

14. Jalili T, Carlstrom J, Kim S, Freeman D, Jin H, Wu TC. Quercetin-supplemented diets lower blood pressure and attenuate cardiac hypertrophy in rats with aortic constriction. *J Cardiovascular Pharmacol* 47(4): 531-541, 2006.

15. Jin F, Nieman DC, Shanely RA, Knab AM, Austin MD, Sha W. The variable plasma quercetin response to 12-week quercetin supplementation in humans. *Eur J Clin Nutr.* 64(7): 692-697, 2010.

16. Joanny, P, Steinberg, J, Robach, P, Richalet, JP, Gortan, C, Gardette, B and Jammes, Y. Operation Everest III (Comex'97): the effect of simulated severe hypobaric hypoxia on lipid peroxidation and antioxidant defense systems in human blood at rest and after maximal exercise. *Resuscitation* 49: 307-314, 2001.

17. Katayama K, Goto K, Ishida K, Ogita F. Substrate utilization during exercise and recovery at moderate altitude. *Metab Clin Exper* 59(7): 959-966, 2009.

18. Khoo NK, White CR, Pozzo-Miller L, Zhou F, Constance C, Inoue T. Dietary flavonoid quercetin stimulates vasorelaxation in aortic vessels. *Free Radical Biol Med* 49(3): 339-347, 2010.

19. Lee KH, Park E, Lee HJ, Kim MO, Cha YJ, Kim JM. Effects of daily quercetin-rich supplementation

on cardiometabolic risks in male smokers. *Nutr Res Practice* 5(1): 28-33, 2011.

20. MacRae HS, Mefferd KM. Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *Int J Sport Nutr Exerc Metab* 16(4): 405-419, 2006.

21. Nieman DC, Williams AS, Shanely RA, Jin F, McAnulty SR, Triplett NT. Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Med Sci Sports Exerc* 42(2): 338-345, 2010.

22. Nicholson SK, Tucker GA, Brameld JM. Physiological concentrations of dietary polyphenols regulate vascular endothelial cell expression of genes important in cardiovascular health. *Br J Nutr* 103(10): 1398-1403, 2009.

23. Punithavathi VR, Stanely Mainzen Prince P. The cardioprotective effects of a combination of quercetin and alpha-tocopherol on isoproterenol-induced myocardial infarcted rats. *J Biochem Mol Toxicol* 25(1): 28-40, 2011.

24. Purushothaman J, Suryakumar G, Shukla D, Malhotra AS, Kasiganesan H, Kumar R. Modulatory effects of seabuckthorn (*Hippophae rhamnoides* L.) in hypobaric hypoxia induced cerebral vascular injury. *Brain Res Bull* 77(5): 246-252, 2008.

25. Subudhi AW, Jacobs KA, Hagobian TA, Fattor JA, Muza SR, Fulco CS, Cymerman A, Friedlander AL. Changes in ventilator threshold at high altitude: effect of antioxidants. *Med Sci Sports Exerc* 38(8): 1425-1431, 2006.

26. Vasquez-Garzon VR, Arellanes-Robledo J, Garcia-Roman R, Aparicio-Rautista DI, Villa-Trevino S. Inhibition of reactive oxygen species and pre-neoplastic lesions by quercetin through an antioxidant defense mechanism. *Free Radical Res* 43(2): 128-137, 2009.

27. West JB, Hackett PH, Maret KH, Milledge JS, Peters RM, Pizzo CJ. Pulmonary gas exchange on the summit of Mount Everest. *J Appl Physiol Resp Environment Exerc Physiol* 55(3): 678-687, 1983.

28. Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of

## QUERCETIN SUPPLEMENTATION AND SIMULATED ALTITUDE PERFORMANCE

93 intervention studies. *Am J Clin Nutr* 81(1 Suppl): 243S-255S, 2005.

29. Wilson RT, Wang J, Chinchilli V, Richie JP, Virtamo J, Moore LE. Fish, vitamin D, and flavonoids in relation to renal cell cancer among smokers. *Am J Epidemiol* 170(6): 717-729, 2009

30. Xu F, Proft J, Gibbs S, Winkfein B, Johnson JN, Syed N. Quercetin targets cysteine string protein (CSPalpha) and impairs synaptic transmission. *PloS one* 5(6): 2010.

31. Zhou J, Zhou S, Gao Y, Zeng S. Modulatory effects of quercetin on hypobaric hypoxic rats. *Eur J Pharmacol* 674(2-3): 450-454, 2011.