

4-2015

Grip Strength and Functional Measures in the Mature Adult: Brief Report II

Mark DeBeliso

Mikaela Boham

Chad Harris

Carole Carson

Joseph M. Berning

See next page for additional authors

Follow this and additional works at: https://digitalcommons.csumb.edu/kin_fac

This Article is brought to you for free and open access by the Kinesiology at Digital Commons @ CSUMB. It has been accepted for inclusion in Kinesiology Faculty Publications and Presentations by an authorized administrator of Digital Commons @ CSUMB. For more information, please contact digitalcommons@csumb.edu.

Authors

Mark DeBeliso, Mikaela Boham, Chad Harris, Carole Carson, Joseph M. Berning, Trish Sevene, Kent J. Adams, and Mike Climstein



Grip Strength and Functional Measures in the Mature Adult: Brief Report II

Mark DeBeliso¹, Mikaela Boham², Chad Harris³, Carole Carson⁴, Joseph M. Berning⁵, Trish Sevene⁶, Kent J. Adams⁷,
Mike Climstein⁸

¹Southern Utah University, Cedar City, Utah, USA

²Texas A&M Corpus Christi, Corpus Christi, Texas, USA

³Central Oregon Community College, Bend, Oregon, USA

^{4,5}New Mexico State University, Las Cruces, New Mexico, USA

^{6,7}California State University Monterey Bay, Seaside, California, USA

⁸University of Sydney, Sydney, AUS

(¹markdebeliso@suu.edu)

Abstract- Grip strength has been shown to be closely linked to body strength in mature aged adults. However what may be of greater importance is the relationship between grip strength and functional movements in aging adults. Purpose: This study examined the relationship between grip strength and functional body movements in mature aged adults. Methods: Female (n=12, age=71.2±3.8 years, mass=66.3±9.2 kg) and male (n=16, age=72.9±4.7 years, mass=85.5±9.4 kg) participants completed functional body movements including: vertical jump (VJ), medicine ball (MB) throws (1.5, 3.0, & 5.0 Kgs), and a stair climb (SC) test. Likewise, all participants performed maximal hand grip (MG) with the Jamar hand grip dynamometer. Pearson correlation coefficients (PCC) were then calculated to determine the relationship between dominant hand MG and the functional body movement scores. Results: PCC's were as follows: MG-VJ (r=0.62), MG-MB^{1.5} (r=0.86), MG-MB^{3.0} (r=0.87), MG-MB^{5.0} (r=0.91), and MG-SCP (r=0.79). All PCC's were statistically significant at P<0.01. Conclusions: Within the parameters of this study, MG strength is strongly reflective of functional body movements that require upper and lower body power output. In this regard, assessing grip strength may be beneficial to clinicians interested in assessing functional body movements in aging adults.

Keywords- grip, grip strength, mature adult

I. INTRODUCTION

Aging is associated with muscle loss and strength a condition referred to as sarcopenia (1). Sarcopenia related loss of muscle strength and muscle function is associated with decrements in the ability to perform activities of daily living (ADLs) (2) and an increased risk of falls (3). Injuries related to falls often lead to permanent disability resulting in loss of independence (4).

Given the association between age related muscle loss/strength and mortality/injuries as well as the loss in the

ability to perform ADLs, it may be of value to develop a practical clinical test that could provide insight into an aging individual's ability to perform ADLs.

In this regard, grip strength assessment may be of value. Consider the following: grip strength has been shown to be associated with reduced mortality in both the young and mature aged adults (5,6). In our previous work (7) we focused upon the relation between grip strength and muscle strength in regions of the body; which confirmed our hypothesis that grip strength is positively related to muscle strength in other regions of the body. Given our previous findings (7) we wanted to explore if there was a positive relationship between grip strength and functional body movements. Given the relationship between grip and muscle strength in other regions of the body, we felt it was reasonable to hypothesize that grip strength would be positively related to functional body movements. Our assumption is that functional body movements are representative of the ability to perform ADLs.

Hence, the purpose of this study was to determine if a meaningful positive relationship existed between grip strength and functional movements in a participant pool of mature aged adults. This brief report is a continuation of our earlier work (7).

II. METHODS

Participants

Participants (n = 28) included both males and females in either their 7th, 8th, or 9th decade of life (68-88 years). Recruitment strategies included public announcements, flyers, and word of mouth. The participants were independent and community-dwelling with no previous background in resistance training. Participants were cleared for participation in the study by their personal physician. Prior to the execution of the study, all participants were verbally informed of the details of the study, read and signed an informed consent

document approved by an Institutional Review Board for the use of Human Subjects.

TABLE I. PARTICIPANT DESCRIPTIVE CHARACTERISTICS (MEAN±SD)

Participants	N	Age (years)	Body Mass (kilograms)
Female	12	71.2±3.8	66.3±9.2
Male	16	72.9±4.7	85.5±9.4

Procedures

Maximal grip strength (MG) was assessed with a Jamar hand dynamometer. Participants completed two trials of maximal grip with both the dominant and non-dominant hand. Participants were seated with the shoulder at 0° abduction and flexion with the elbow at 90° flexion, as recommended by American Society of Hand Therapists (8). Participants were instructed to familiarize themselves with the Jamar by holding and squeezing the device prior to performing the maximal grip trials. Participants were then instructed to squeeze the device with a maximal effort for three seconds. The trials were separated by approximately 1 minute rest periods. The greatest MG score from the two trials was used for analysis.

Vertical jump (VJ) measures were collected with the Vertec vertical jump measuring device in a manner previously described (9). The participants were given three VJ attempts with a self-selected rest period between trials ranging from 1-3 minutes. The best score (centimeters) was recorded and used for subsequent analysis. Vertical jump measures collected in this manner have been reported as reliable with intra class correlations (ICC's) ranging from 0.87-0.89 (9).

Medicine ball (MB) throws were collected for three different MB masses: 1.5, 3.0, and 5.0 Kgs. The MB throws were collected (meters) with the participants in a seated position as previously described (10). The participants were given three trials at each of the three MBs. The participants were allowed a self-selected rest period between trials ranging from 1-3 minutes. Medicine ball measures collected in this manner have been reported as reliable with ICC's ≥0.98 (10).

Stair climb (SC) measures were collected for each participant as previously described (11). The best score of two trials separated by a 3 minute rest period was used for subsequent analysis. There were 12 stairs at 15.24 cm in height for a total of 1.83 meters.

Prior to study initiation, participants were instructed in proper execution of each task and appropriate breathing patterns in order to minimize cardiovascular stress (12). The participants were also allowed to practice each task assuring that they were familiar with the body mechanics of each movement.

Analysis

A personal computer with Microsoft Excel 2013 software was utilized for data management and statistical analysis. Standard descriptive statistics (mean and standard deviation) for age, height, and body mass were calculated. Pearson

correlation coefficients (PCC) were calculated between maximal dominant grip strength and functional movement measures (significance $\alpha \leq 0.05$).

III. RESULTS

All of the participants were able to complete the MG and five functional body movement assessments (MG, VJ, MB^{1.5}, MB^{3.0}, MB^{5.0}, and SC). Table 1 provides the subject descriptive statistics for age and body mass (mean± standard deviation).

Table 2 provides the Pearson correlation coefficients (r) between MG and the functional body movement scores. The PCC's ranged from high (r=0.62) to very high (r=0.91) (13). All PCC's were statistically significant (p<0.01).

TABLE II. PEARSON CORRELATION COEFFICIENTS (R) BETWEEN MAXIMAL DOMINANT GRIP STRENGTH AND FUNCTIONAL BODY MOVEMENTS

	Vertical Jump	Medicine Ball 1.5 Kg	Medicine Ball 3.0 Kg	Medicine Ball 5.0 Kg	Stair Climb
Grip Strength	0.62*	0.86*	0.87*	0.91*	-0.37*

*p<0.01

TABLE III. FUNCTIONAL BODY MOVEMENTS SCORES (MEAN±SD)

Strength Measure	Male (n=16)	Female (n=12)
Maximal Grip (Kg)	30.3±5.6	10.6±3.3
Vertical Jump (cm)	27.9±6.8	18.7±5.3
Medicine Ball 1.5 Kg (m)	5.1±0.7	3.4±0.5
3.0 Kg (m)	4.0±0.5	2.7±0.4
5.0 Kg (m)	3.3±0.3	2.2±0.3
Stair Climb (seconds)	2.9±0.5	3.3±0.6

Table 3 lists the MG and functional body movement scores for all of the variables collected for both genders (mean±SD).

IV. DISCUSSION

The purpose of this study was to determine if a meaningful positive relationship existed between maximal grip strength and functional body movements in mature adults. The relationships between MG and the functional body movements in this study ranged from high to very high with exception of the timed SC.

The functional body movements (VJ, MB and SC) selected for this study were all muscular power oriented for both the upper and lower body. When the SC scores were converted to measures of power output (Watts=Newtons*meters/second) the PCC between MG and stair climb power output (SCP) was also high (r=0.79). Hence, the PCC's reported in this study as a

measure of muscular power output ranged from high to very high (13).

As previously stated (7), the MGs recorded during the current study were inexplicably lower than norms reported elsewhere (14). VJ measures were consistent with those reported by Perchthaler et al. (15). The MB throws recorded in this study were also consistent with previously reported MB scores in the mature adult population (10). The SC times were also very close to those in a previous study using a similar participant pool and stair case (step number and height) (11).

A previous meta-analysis revealed that the odds ratios for mortality as a function of grip strength persisted after correcting for gender (6). As such, we included both genders in the calculation of the PCCs.

During the development of this study we aimed to select functional body movements that required muscular power. While our previous work established a strong link between MG and body strength in mature adults (7), we felt that muscular power, the combination of strength and the rate that it can be applied may be more important with regards to the ability to perform ADLs. Research indicates that the ability to develop a high degree of muscular power appears to diminish with age (16,17). An increase in power enables the older adult to improve performance in tasks that require strength with speed, which relates to many functional and recreational activities (climbing stairs, walking/gait speed, recovering from a fall, golf, biking, etc.) (17,18). For example, Bassey et al. (1992) found strong correlations between leg extensor power and functional abilities such as chair-rising speed, stair-climbing speed, walking speed, and stair climbing power (19).

Consistent with our previous work (7), the results of our current study demonstrate that MG is equally effective with regards to estimating functional body movements that require the application of muscular power.

A limitation to the current study is the number of participants. Future studies examining MG and estimating functional body movements should grow the participant pool in order to verify the relationships established in the current study. It would also be of interest to determine if subsequent exercise prescription that is based on low MG leads to advanced physical capabilities as is seen with those participating in masters caliber sports (20,21,22,23,24,25).

Exercise prescription focusing on improving muscular strength in mature adults has proven effective (26,27,28). With that said, clinicians would be well advised to include exercises that mimic ADLs in to any exercise prescription. This process would best assure that muscular strength gains as a result of implementing an exercise protocol would translate into improved functionality. This process would be analogous to athletes including skill specific training and exercise modalities that focus on specificity of training in order to optimize a transfer of muscular strength and power to the field of play (29).

Additional future studies should also examine different aspects of MG as found with advanced technology such as the Grip Force Map system (30,31,32). The Grip Force Map

system employs tactile array technology that allows a detailed force map of the hand while gripping an optimal diametered cylinder that is covered with a tactile array surface. The force map provides information regarding the force and pressure generated by each finger within the context of the entire hand. Detailed information regarding MG may provide further insight as the relationship between MG and functional body movements.

Given the high PCCs between MG and functional body movements established in this study, it appears that grip strength measures may serve as a clinically friendly screening assessment of functional body movements. Grip strength measures could be easily assessed as part of an annual physical exam allowing practitioners a starting point for exercise prescription. Such a practice would be of particular value to patients experiencing sarcopenia.

V. CONCLUSIONS

Within the parameters of this study it is concluded that:

1. Maximal dominant grip strength is positively associated with functional body movements that are muscular power oriented for both the upper and lower body.
2. Given the relationship between MG and functional body movements, clinicians might be well advised to collect MG as part of a mature adult's annual exam.

ACKNOWLEDGMENT

To our friend and mentor, Doctor John Patrick O'Shea, you're missed and never forgotten.

REFERENCES

- [1] Cruz-Jentoft, A.J., Baeyens, J.P., Bauer, J.M., et al. (2010). Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*, 39, 412-423.
- [2] Bastiaanse, L.P., Hilgenkamp, T.I., Echteid, M.A., & Evenhuis, H.M. (2012). Prevalence and associated factors of sarcopenia in older adults with intellectual disabilities. *Research in Developmental Disabilities*, 33, 2004-2012.
- [3] Landi F, Liperoti R, Fusco D, et al. (2012). Prevalence and risk factors of sarcopenia among nursing home older residents. *Journal of Gerontology: Biological Sciences*, 67, 48-55.
- [4] World Health Organization. (2007). WHO global report on falls prevention in older age. WHO Press, World Health Organization, Geneva, Switzerland.
- [5] Ruiz, J.R., Sul, X., Lobelo, F., Morrow, J.R., Jackson, A.W., Sjostrom, M., & Blair, S.N. (2008). Association between muscular strength and mortality in men: prospective cohort study. *British Medical Journal*, 337:a439.
- [6] Cooper, R., Kuh, D., & Hardy, R. (2010). Objective measured physical capability levels and mortality: systematic review and meta-analysis. *British Medical Journal*, 341:C4467.

- [7] DeBeliso, M., Boham, M., Carson, C., Harris, C., Berning, J.M., Sevene, T., & Adams, K.J. (2015). Grip and body strength measures in the mature adult: A Brief Report. *International Journal of Science and Engineering Investigations*, 4(37), 83-86.
- [8] Fess, E.E. Grip strength. In: Casanova, J.S. (Ed.) *Clinical assessment recommendations*, 2nd Edn. Chicago, American Society of Hand Therapists, 1992, 41-45.
- [9] Nuzzo, J.L., Anning, J.H., & Scharfenberg, J.M. (2011). The reliability of three devices used for measuring vertical jump height. *Journal of Strength & Conditioning Research*, 25(9), 2580-2590.
- [10] Harris, C., Wattles, A.P., DeBeliso, M., Sevene-Adams, P.G., Berning, J.M., & Adams, K.J. (2011). The seated medicine ball throw as a test of upper body power in older adults. *Journal of Strength and Conditioning Research*, 25(8), 2344.
- [11] Hanson, E.D., Srivatsan, S.R., Agrawal, S., Menon, K.S., Delmonico, M.J., Wang, M.Q., & Hurley, B.F. (2009). Effects of strength training on physical function: influence of power, strength, and body composition. *Journal of Strength Conditioning Research*, 23(9), 2627-2637.
- [12] Evans, W. (1999). Exercise guidelines for the elderly. *Medicine & Science in Sports & Exercise*, 31(1), 12-17.
- [13] Safrit, M.J., & Wood, T.M. (1995). *Introduction to measurement in physical education and exercise* (3rd Ed.) St. Louis, MO: Mosby.
- [14] Bohannon, R., Peolsson, A., Massy-Westropp, N., Desrosiers, J., & Bear-Lehman, J. (2006). Reference values for adult grip strength measured with a Jamar dynamometer: a descriptive meta-analysis. *Physiotherapy*, 92(1), 11-15.
- [15] Perchthaler, D., Grau, S., & Hein, T. (2015) Evaluation of a 6-week whole-body vibration intervention on neuromuscular performance in older adults. *Journal of Strength Conditioning Research*, 29(1), 86-95.
- [16] Brill, P. A., C. A. Macera, D. R. Davis, S. N. Blair, and N. Gordon. (2000). Muscular strength and physical function. *Medicine and Science in Sports and Exercise*, 32, 412-416.
- [17] Riley, P. O., M. L. Schenkman, R. W. Mann, et al. (1991). Mechanics of a constrained chair-rise. *Journal of Biomechanics*, 24, 77-85.
- [18] McFaden, B.J., & Winter, D.A. (1988). An integrated biomechanical analysis of normal stair ascent. *Journal of Biomechanics*, 24, 733-744.
- [19] Bassey, E.J., Fiatarone, M.A., O'Neill, E.F., Kelly, M., Evans, W.J., and Lipsitz, L.A. (1992). Leg extensor power and functional performance in very old men and women. *Clinical Science*, 82, 321-327.
- [20] DeBeliso, M., Sevene, T.G., Walsh, J., Adams, K.J., Kettunen, J., Heazlewood, I.T., & Climstein, M. (2014). Body mass index of north american participants at the World Masters Games. *Journal of Sport Science*, 2, 189-194.
- [21] DeBeliso, M., Walsh, J., Climstein, M., Heazlewood, I.T., Kettunen, J., Sevene, T.G., & Adams, K.J. (2014). World Masters Games: North american participant medical and health history survey. *The Sport Journal*, 0417.
- [22] Walsh, J., Climstein, M., Heazlewood, I.T., DeBeliso, M., et al. (2013). Masters Athletes: No evidence of increased incidence of injury in football code athletes. *Advances in Physical Education*, 3(1), 36-42.
- [23] Heazlewood, I.T., Walsh, J., Climstein, M., Adams, K.J., DeBeliso, M., et al. (2013). Gender differences in motivating participation and adherence in masters sport. *Journal of Science and Medicine in Sport*, 16(6), S115.
- [24] Walsh, J., Climstein, M., Heazlewood, I.T., Kettunen, J., DeBeliso, M., & Adams, K.J. (2013). Body mass index for athletes participating in swimming at the World Masters Games. *Journal of Sports Medicine and Physical Fitness*, 53, 162-168.
- [25] Heazlewood, I.T., Walsh, J., Climstein, M., Adams, K.J., Sevene, T.G., DeBeliso, M., & Kettunen, J. (2013). Injuries of athletes in training for 2010 Pan Pacific Masters Games: Types and location. *Journal of Science and Medicine in Sport*, 16(6), S54.
- [26] Harris, C., DeBeliso, M., Spitzer-Gibson, T., & Adams, K. (2004). The effect of resistance training intensity on strength gain response in the older adult. *Journal of Strength and Conditioning Research*, 18(4): 833-838.
- [27] DeBeliso, M., Harris, C., Spitzer-Gibson, T., & Adams, K. (2005). A comparison of periodized and fixed repetition training protocols on strength measures in older adults. *Journal of Science & Medicine in Sport*, 8(2): 190-199.
- [28] Harris, C., DeBeliso, M., Adams, K., Irmischer, B. & Gibson, T.A. (2007). Detraining in the older adult: effects of prior training intensity on strength retention. *Journal of Strength and Conditioning Research*, 21(3): 813-818.
- [29] O'Shea, J. P. (200). *Quantum Strength & Power Training II: Gaining the Winning Edge*. Patrick's Books, Corvallis, OR.
- [30] DeBeliso, M., Fichter, D., Murdock, L., & Adams, K.J. (2009). Grip norms and reliability of the hand grip ForceMap system. *Medicine & Science in Sports & Exercise*, 41(5), S430.
- [31] DeBeliso, M., McChesney, J.W., & Murdock, L.E. (2013). Grip norms and reliability of the hand grip ForceMap system. *Journal of Hand Surgery*, 38(9), 1009-10.
- [32] McChesney, J.W., DeBeliso, M., & Murdock, L.E. (2012). The hand grip ForceMap system. *International Journal of Science and Engineering Investigations*, 1(8), 76-78.

Mark DeBeliso, PhD is a Professor and Graduate Program Director of the Masters of Science in Sport Conditioning and Performance at Southern Utah University, Utah, USA. His research interests include mechanics and metabolics of sport movements and work tasks, strength training for all walks of life, orthopedic biomechanics, and masters athletes.

Mikaela Boham, EdD is an Assistant Professor and Director of the Athletic Training Program at Texas A&M University Corpus Christi, Texas, USA. Her research interests include sports injuries and traumatic brain injuries in sports.

Carole Carson, MS is the Director of Nursing at the Rehabilitation Hospital of Southern New Mexico, Las Cruces, New Mexico, USA. She also serves as an Adjunct Lecturer for New Mexico State University with research interests in strength and rehabilitation.

Chad Harris, PhD is the Instructional Dean of Health Science at Central Oregon Community College, Oregon, USA. His research interests include training effects on power production, weightlifting biomechanics, senior strength training and metabolic responses to power training.

Joseph M. Berning, PhD is a Professor and Director of the Exercise Physiology Lab in the Department of Human Performance, Dance & Recreation at New Mexico State University, Las Cruces, New Mexico, USA. His research interests include strength and power training, overtraining, and warm-up strategies to enhance performance.

Trish Sevene, PhD is an Associate Professor in the Kinesiology Department at California State University Monterey Bay, California, USA. Her research interests include the biological basis of human performance and aging, work-related lifting tasks, and masters athletes.

Kent J. Adams, PhD is a Professor and Chair of the Kinesiology Department at California State University Monterey Bay, California, USA. His research interests include strength and power training across the lifespan, work-related lifting tasks, and masters athletes.

Mike Climstein, PhD is an Honorary Associate Professor with the Health & Performance Faculty Research Group at the University of Sydney, AUS. His research interests include medical and health history of Masters Games and World Masters games athletes.